

The braided monoidal structure on the category of Hom-type Doi-Hopf modules

Daowei Lu*, Shuanhong Wang

Abstract. Let (H, α_H) be a Hom-Hopf algebra, (A, α_A) a right H -comodule algebra and (C, α_C) a left H -module coalgebra. Then we have the category ${}_A\mathcal{M}(H)^C$ of Hom-type Doi-Hopf modules. The aim of this paper is to make the category ${}_A\mathcal{M}(H)^C$ into a braided monoidal category. Our construction unifies quasitriangular and coquasitriangular Hom-Hopf algebras and Hom-Yetter-Drinfeld modules. We study tensor identities for monoidal categories of Hom-type Doi-Hopf modules. Finally we show that the category ${}_A\mathcal{M}(H)^C$ is isomorphic to $A\#C^*$ -module category.

Keywords: Hom-Hopf algebra; Drinfeld double; Braided monoidal category; Doi-Hopf module.

Mathematics Subject Classification: 16W30, 16T05.

Introduction

The Doi-Hopf datum (H, A, C) introduced in [5] consists of a Hopf algebra H , a right H -comodule algebra A and a left H -module coalgebra C . The Doi-Hopf module M over (H, A, C) is both a left A -module and a right C -comodule satisfying certain compatible condition. The category of Doi-Hopf modules over (H, A, C) is denoted by ${}_A\mathcal{M}(H)^C$. The research of ${}_A\mathcal{M}(H)^C$ turn out to be very essential: it is pointed out in [5] that categories such as module and comodule over bialgebra, in [13] that the Hopf modules category, and in [3, 12] that the Yetter-Drinfeld modules category are special cases of ${}_A\mathcal{M}(H)^C$. For a further study of Doi-Hopf modules, we refer to [4].

Braided monoidal categories give rise to solutions to the Quantum Yang-Baxter equations. The classical braided monoidal categories come from the representations of quasitriangular Hopf algebras which have been very widely studied. One of the most important examples of quasitriangular Hopf algebras is the Drinfeld double of any finite dimensional

*Corresponding author

Hopf algebra. As such, they are interesting to different research communities in mathematical physics (see [8] for example).

From a physical viewpoint, Hom-type structures are important because they are related to vertex operator algebras and string theory (see [1],[2]). Hom-Lie algebras were introduced in [6] to describe the structures on some q -deformations of the Witt and the Virasoro algebras, both of which are important in vertex operator algebras and string theory. Also in this paper Hom-type algebras have been introduced in the form of Hom-Lie algebras, where the Jacobi identity was twisted along a linear endomorphism. Meanwhile, Hom-associative algebras have been suggested in [10] to give rise to a Hom-Lie algebra using the commutator bracket. Other Hom-type structures such as Hom-coalgebras, Hom-bialgebras, Hom-Hopf algebras as well as their properties have been considered in [11].

Yang-Baxter equations and its solutions have been generalized to Hom-bialgebra in [14], and the author pointed out that a quasitriangular Hom-bialgebra (H, R) provides solutions of a class of Hom Yang-Baxter equations:

$$\begin{aligned}(R_{12}R_{13})R_{23} &= R_{23}(R_{13}R_{12}), \\ R_{12}(R_{13}R_{23}) &= (R_{23}R_{13})R_{12}.\end{aligned}$$

Motivated by these ideas, in this paper our aim is to make the Hom-type Doi-Hopf module category ${}_A\mathcal{M}(H)^C$ into a braided monoidal category. We will prove that ${}_A\mathcal{M}(H)^C$ and ${}_{A\#C^*}\mathcal{M}$ are isomorphic as braided monoidal category. Hence $A\#C^*$, as a generalization of the Drinfeld double $D(H)$, is quasitriangular.

This article is organized as follows:

In section 1, we will recall the basic definitions and results on Hom-Hopf algebra, including Hom-Hopf algebra, Hom-module coalgebra, Hom-comodule algebra and quasitriangular Hom-bialgebra.

In section 2, we will discuss when the category ${}_A\mathcal{M}(H)^C$ becomes monoidal (see Proposition 2.3). Then Yetter-Drinfeld category ${}_H\mathcal{YD}^H$ is isomorphic to ${}_H\mathcal{M}(H^{op} \otimes H)^H$ as monoidal category. Maps between the underlying Hom-Hopf algebras, Hom-algebras and Hom-coalgebras give rise to functors between Hom-type Doi-Hopf modules (see Proposition 2.6 and 2.7). We will also discuss particular situations which give rise to the tensor identities (see Theorem 2.8 and 2.10).

In section 3, we will present the necessary and sufficient conditions for ${}_A\mathcal{M}(H)^C$ to be braided. We point out that this comes down to a convolution invertible map $Q : C \otimes C \rightarrow A \otimes A$ subject to certain compatible relations (see Theorem 3.8). Especially, on one hand when $C = k$ this conditions turn out to be equivalent to $(A, Q(1))$ being quasitriangular. On the other hand when $A = k$ then we recover the definition of a coquasitriangular Hom-Hopf algebra.

In section 4, we will prove the monoidal category isomorphism ${}_A\mathcal{M}(H)^C \simeq {}_{A\#C^*}\mathcal{M}$

(see Proposition 4.4). Since ${}_A\mathcal{M}(H)^C$ is braided, so is ${}_A\#C^*\mathcal{M}$. In the case of ${}_H\mathcal{M}(H^{op} \otimes H)^H$, ${}_A\#C^*$ reduces to the Drinfeld double $D(H) = H \bowtie H^*$.

Throughout this article, all the vector spaces, tensor product and homomorphisms are over a fixed field k . For a coalgebra C , we will use the Heyneman-Sweedler's notation $\Delta(c) = c_1 \otimes c_2$, for all $c \in C$ (summation omitted).

1 Preliminary

In this section, we will recall from [9, 11] the basic definitions and results on the Hom-Hopf algebras, Hom-modules and Hom-comodules.

A unital Hom-associative algebra is a triple (A, m, α) where $\alpha : A \longrightarrow A$ and $m : A \otimes A \longrightarrow A$ are linear maps, with notation $m(a \otimes b) = ab$ such that for any $a, b, c \in A$,

$$\begin{aligned}\alpha(ab) &= \alpha(a)\alpha(b), \quad \alpha(1_A) = 1_A, \\ 1_A a &= \alpha(a) = a 1_A, \quad \alpha(a)(bc) = (ab)\alpha(c).\end{aligned}$$

A linear map $f : (A, \mu_A, \alpha_A) \longrightarrow (B, \mu_B, \alpha_B)$ is called a morphism of Hom-associative algebra if $\alpha_B \circ f = f \circ \alpha_A$, $f(1_A) = 1_B$ and $f \circ \mu_A = \mu_B \circ (f \otimes f)$.

A counital Hom-coassociative coalgebra is a triple $(C, \Delta, \varepsilon, \alpha)$ where $\alpha : C \longrightarrow C$, $\varepsilon : C \longrightarrow k$, and $\Delta : C \longrightarrow C \otimes C$ are linear maps such that

$$\begin{aligned}\varepsilon \circ \alpha &= \varepsilon, \quad (\alpha \otimes \alpha) \circ \Delta = \Delta \circ \alpha, \\ (\varepsilon \otimes id) \circ \Delta &= \alpha = (id \otimes \varepsilon) \circ \Delta, \\ (\Delta \otimes \alpha) \circ \Delta &= (\alpha \otimes \Delta) \circ \Delta.\end{aligned}$$

A linear map $f : (C, \Delta_C, \alpha_C) \longrightarrow (D, \Delta_D, \alpha_D)$ is called a morphism of Hom-coassociative coalgebra if $\alpha_D \circ f = f \circ \alpha_C$, $\varepsilon_D \circ f = \varepsilon_C$ and $\Delta_D \circ f = (f \otimes f) \circ \Delta_C$.

In what follows, we will always assume all Hom-algebras are unital and Hom-coalgebras are counital.

A Hom-bialgebra is a quadruple (H, μ, Δ, α) , where (H, μ, α) is a Hom-associative algebra and (H, Δ, α) is a Hom-coassociative coalgebra such that Δ and ε are morphisms of Hom-associative algebra.

A Hom-Hopf algebra (H, μ, Δ, α) is a Hom-bialgebra H with a linear map $S : H \longrightarrow H$ (called antipode) such that

$$\begin{aligned}S \circ \alpha &= \alpha \circ S, \\ S(h_1)h_2 &= h_1 S(h_2) = \varepsilon(h)1,\end{aligned}$$

for any $h \in H$. For S we have the following properties:

$$\begin{aligned} S(h)_1 \otimes S(h)_2 &= S(h_2) \otimes S(h_1), \\ S(gh) &= S(h)S(g), \quad \varepsilon \circ S = \varepsilon. \end{aligned}$$

For any Hopf algebra H and any Hopf algebra endomorphism α of H , there exists a Hom-Hopf algebra $H_\alpha = (H, \alpha \circ \mu, 1_H, \Delta \circ \alpha, \varepsilon, S, \alpha)$.

Let (A, α_A) be a Hom-associative algebra, M a linear space and $\alpha_M : M \longrightarrow M$ a linear map. A left A -module structure on (M, α_M) consists of a linear map $A \otimes M \longrightarrow M$, $a \otimes m \mapsto a \cdot m$, such that

$$\begin{aligned} 1_A \cdot m &= \alpha_M(m), \\ \alpha_M(a \cdot m) &= \alpha_A(a) \cdot \alpha_M(m), \\ \alpha_A(a) \cdot (b \cdot m) &= (ab) \cdot \alpha_M(m), \end{aligned}$$

for any $a, b \in A$ and $m \in M$.

Similarly we can define the right A -modules. Let (M, μ) and (M', μ') be two left A -modules, then a linear map $f : M \rightarrow N$ is called left A -module map if $f(a \cdot m) = aa \cdot f(m)$ for any $a \in A$, $m \in M$ and $f \circ \mu = \mu' \circ f$.

Suppose that (M, μ) is a right A -module and (N, ν) is a left A -module, then denote by X the subspace of $M \otimes N$ spanned by the elements of the type $\left\{ \sum m_i \cdot a \otimes \nu(n_i) - \sum \mu(m_i) \otimes a \cdot n_i \mid \forall m_i \in M, n_i \in N, a \in A \right\}$. Then we have the quotient space

$$M \otimes_A N = (M \otimes N)/X.$$

Let (C, α_C) be a Hom-coassociative coalgebra, M a linear space and $\alpha_M : M \longrightarrow M$ a linear map. A right C -comodule structure on (M, α_M) consists a linear map $\rho : M \longrightarrow M \otimes C$ such that

$$\begin{aligned} (id \otimes \varepsilon_C) \circ \rho &= \alpha_M, \\ (\alpha_M \otimes \alpha_C) \circ \rho &= \rho \circ \alpha_M, \\ (\rho \otimes \alpha_C) \circ \rho &= (\alpha_M \otimes \Delta) \circ \rho. \end{aligned}$$

Let (M, μ) and (M', μ') be two right (C, γ) -comodules, then a linear map $g : M \longrightarrow M'$ is called right C -comodule map if $g \circ \mu = \mu' \circ g$ and $\rho_{M'} \circ g = (g \otimes id) \circ \rho_M$.

Suppose that (M, μ) is a right C -comodule and (N, ν) is a left C -comodule, then we have the cotensor product

$$M \square_C N = \left\{ \sum m_i \otimes n_i \in M \otimes N \mid \sum \mu(m_i) \otimes n_{i(-1)} \otimes n_{i(0)} = \sum m_{i(0)} \otimes m_{i(1)} \otimes \nu(n_i) \right\}.$$

Let (H, α_H) be a Hom-bialgebra. A Hom-algebra (A, α_A) is called a right H -comodule algebra if A is a right H -comodule via ρ , and

$$\rho(ab) = \rho(a)\rho(b), \quad \rho(1_A) = 1_A \otimes 1_H,$$

for all $a, b \in A$. A Hom-coalgebra (C, α_C) is called a left H -module coalgebra if C is a left H -module and

$$\Delta_C(h \cdot c) = \Delta_H(h) \cdot \Delta_C(c), \quad \varepsilon_C(h \cdot c) = \varepsilon_H(h)\varepsilon_C(c),$$

for all $h \in H, c \in C$.

Suppose that (A, α_A) is a right H -comodule algebra and (C, α_C) is a left H -module coalgebra. An object (M, α_M) is called a Doi-Hopf module over (H, A, C) if M is a left A -module via $\cdot$ and a right C -comodule via ρ' satisfying

$$\rho(a \cdot m) = a_{(0)} \cdot m_{(0)} \otimes a_{(1)} \cdot m_{(1)},$$

for all $a \in A$ and $m \in M$.

The category of Doi-Hopf modules over (H, A, C) will be denoted by ${}_A\mathcal{M}(H)^C$.

Recall from [7] that a monoidal category $(\mathcal{C}, \otimes, I, a, l, r)$ is braided if there exists a family of natural isomorphisms $t = \{t_{V,W} : V \otimes W \rightarrow W \otimes V\}_{V,W \in \mathcal{C}}$ such that

$$a_{V,W,U} t_{U,V \otimes W} a_{U,V,W} = (id_V \otimes t_{U,W}) a_{V,U,W} (t_{U,V} \otimes id_W), \quad (1.1)$$

$$a_{W,U,V}^{-1} t_{U \otimes V,W} a_{U,V,W}^{-1} = (t_{U,W} \otimes id_V) a_{U,W,V}^{-1} (id_U \otimes t_{V,W}), \quad (1.2)$$

for all objects U, V, W in $\mathcal{C}$.

Hom-bialgebra (H, α_H) is quasitriangular [14] if there exists an element $R = R^1 \otimes R^2 \in H \otimes H$ such that

$$\begin{aligned} \Delta^{op}(h)R &= R\Delta(h), \\ (\Delta \otimes \alpha_H)R &= R^{13}R^{23}, \\ (\alpha_H \otimes \Delta)R &= R^{13}R^{12}, \end{aligned}$$

where $R^{13} = R^1 \otimes 1 \otimes R^2$, $R^{12} = R^1 \otimes R^2 \otimes 1$ and $R^{23} = 1 \otimes R^1 \otimes R^2$.

2 Monoidal structure on the category ${}_A\mathcal{M}(H)^C$

In this section, we will discuss when the category ${}_A\mathcal{M}(H)^C$ becomes monoidal. In what follows, let (H, α_H) be a Hom-Hopf algebra, (A, α_A) a right H -comodule algebra and (C, α_C) a left H -module coalgebra. First of all, we have the following definition.

Definition 2.1. The Doi-Hopf datum (H, A, C) called is a monoidal Doi-Hopf datum if (A, α_A) and (C, α_C) are Hom-bialgebras with the following compatible conditions:

$$a_{(0)1} \otimes a_{(0)2} \otimes \alpha_H^2(a_{(1)}) \cdot (cd) = a_{1(0)} \otimes a_{2(0)} \otimes (a_{1(1)} \cdot c)(a_{2(1)} \cdot d), \quad (2.1)$$

$$\varepsilon(a)1_C = \varepsilon(a_{(0)})a_{(1)} \cdot 1_C. \quad (2.2)$$

Remark 2.2. If (H, A, C) is a monoidal Doi-Hopf datum, then A and C are also Doi-Hopf modules. The left A -action and the right C -coaction on C are given by

$$a \cdot c = \varepsilon(a_{(0)})a_{(1)} \cdot c, \quad \rho(c) = c_1 \otimes \alpha_C^{-1}(c_2).$$

The left A -action and the right C -coaction on A are given by

$$a \cdot b = \alpha_A^{-1}(a)b, \quad \rho(a) = a_{(0)} \otimes a_{(1)} \cdot 1_C.$$

Proposition 2.3. Let (H, A, C) be a monoidal Doi-Hopf datum. For all Doi-Hopf modules (M, α_M) and (N, α_N) , $(M \otimes N, \alpha_M \otimes \alpha_N)$ still makes a Doi-Hopf module with the following structures:

$$a \cdot (m \otimes n) = a_1 \cdot m \otimes a_2 \cdot n, \quad (2.3)$$

$$\rho(m \otimes n) = m_{(0)} \otimes n_{(0)} \otimes \alpha_C^{-2}(m_{(1)}n_{(1)}), \quad (2.4)$$

for all $a \in A, m \in M$ and $n \in N$. The category $\mathcal{C} = {}_A\mathcal{M}(H)^C$ is a monoidal category.

Proof. We need only to verify the compatible condition. Indeed for $a \in A, m \in M$ and $n \in N$,

$$\begin{aligned} \rho(a \cdot (m \otimes n)) &= a_{1(0)} \cdot m_{(0)} \otimes a_{2(0)} \cdot n_{(0)} \otimes \alpha_C^{-2}((a_{1(1)} \cdot m_{(1)})(a_{2(1)} \cdot n_{(1)})) \\ &\stackrel{(2.1)}{=} a_{(0)1} \cdot m_{(0)} \otimes a_{(0)2} \cdot n_{(0)} \otimes a_{(1)} \cdot \alpha_C^{-2}(m_{(1)}n_{(1)}) \\ &= a_{(0)} \cdot (m_{(0)} \otimes n_{(0)}) \otimes a_{(1)} \cdot \alpha_C^{-2}(m_{(1)}n_{(1)}). \end{aligned}$$

It is now obvious that tensor product defines a functor $\mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$. Under the trivial A -action and C -coaction, k is a Doi-Hopf module, and k is the unit object in $\mathcal{C}$. Let M, N, P be Doi-Hopf modules, from [9] the isomorphisms

$$\begin{aligned} a_{M,N,P} : (M \hat{\otimes} N) \hat{\otimes} P &\rightarrow M \hat{\otimes} (N \hat{\otimes} P), \quad (m \otimes n) \otimes p \mapsto \alpha_M^{-1}(m) \otimes (n \otimes \alpha_N(p)), \\ l_M : k \hat{\otimes} M &\rightarrow M, \quad \lambda \otimes m \mapsto \lambda \alpha_M^{-1}(m), \\ r_M : M \hat{\otimes} k &\rightarrow M, \quad m \otimes \lambda \mapsto \lambda \alpha_M^{-1}(m), \end{aligned}$$

define respectively the associator, left and right unit of $\mathcal{C}$. □

Example 2.4. Suppose that C is a left H -module bialgebra, which means that C as a Hom-bialgebra is not only a left H -module algebra but also a left H -comodule algebra. Let A be a Hom-bialgebra and a right H -comodule algebra such that for all $a \in A$,

$$a_{(0)1} \otimes a_{(0)2} \otimes a_{(1)1} \otimes a_{(1)2} = a_{1(0)} \otimes a_{2(0)} \otimes a_{1(1)} \otimes a_{2(1)}.$$

Clearly (H, A, C) is a monoidal Doi-Hopf datum.

Actually (H, α_H) itself is a left H -module bialgebra, with the left adjoint action $h \cdot g = (h_1 \alpha_H^{-1}(g)) S \alpha_H(h_2)$ and the coaction afforded by the comultiplication.

Let (H, α_H) be a cocommutative Hom-Hopf algebra, and (C, α_C) be a left H -module bialgebra. Easy to see that (H, H, C) is a monoidal Doi-Hopf datum where H is considered as a right H -comodule via comultiplication.

Proposition 2.5. Let (H, α_H) be a Hom-Hopf algebra with bijective antipode, then H is a right $H^{op} \otimes H$ -comodule with the following structure

$$\rho(h) = \alpha_H^{-1}(h_{12}) \otimes S^{-1} \alpha_H^{-2}(h_{11}) \otimes \alpha_H^{-1}(h_2),$$

for all $h \in H$. Define the left $H^{op} \otimes H$ -action on H by $(h \otimes g) \cdot l = (g \alpha_H^{-1}(l)) \alpha_H(h)$, then $(H^{op} \otimes H, H, H)$ is a Doi-Hopf datum. Moreover we have the category isomorphism

$${}_H \mathcal{M}(H^{op} \otimes H)^H \simeq {}_H \mathcal{YD}^H.$$

Proof. Apparently H is a Doi-Hopf module over the datum $(H^{op} \otimes H, H, H)$. Next we will show that ${}_H \mathcal{M}(H^{op} \otimes H)^H$ is a monoidal category. Take $A = H$ and $C = H^{op}$ as Hom-bialgebra, for all $h, g, g' \in H$,

$$\begin{aligned} & h_{(0)1} \otimes h_{(0)2} \otimes \alpha_H^2(h_1) \cdot (g \bullet g') \\ &= \alpha_H^{-1}(h_{121}) \otimes \alpha_H^{-1}(h_{122}) \otimes (\alpha_H(h_2) \alpha_H^{-1}(g'g)) S^{-1} \alpha_H(h_{11}) \\ &= h_{12} \otimes h_{21} \otimes (h_{22} \alpha_H^{-1}(g'g)) S^{-1} \alpha_H(h_{11}) \\ &= h_{12} \otimes h_{21} \otimes (h_{22} g') (g S^{-1}(h_{11})) \\ &= \alpha_H^{-1}(h_{112}) \otimes \alpha_H^{-1}(h_{212}) \otimes [\alpha_H^{-1}(h_{22} g') (S^{-1}(\alpha_H^{-2}(h_{211})) \alpha_H^{-1}(h_{12}))] (g S^{-1} \alpha_H^{-1}(h_{111})) \\ &= \alpha_H^{-1}(h_{112}) \otimes \alpha_H^{-1}(h_{212}) \otimes [\alpha_H^{-1}(h_{22} g') S^{-1}(\alpha_H^{-1}(h_{211}))] [h_{12} (\alpha_H^{-1}(g) S^{-1} \alpha_H^{-2}(h_{111}))] \\ &= \alpha_H^{-1}(h_{112}) \otimes \alpha_H^{-1}(h_{212}) \\ &\quad \otimes [(S^{-1} \alpha_H^{-2}(h_{111}) \otimes \alpha_H^{-1}(h_{12})) \cdot g] \bullet [(S^{-1}(\alpha_H^{-2}(h_{211}) \otimes \alpha_H^{-1}(h_{22}))) \cdot g'] \\ &= h_{1(0)} \otimes h_{2(0)} \otimes (h_{1(1)} \cdot g) \bullet (h_{1(2)} \cdot g'), \end{aligned}$$

and

$$\varepsilon(h_{(0)}) h_{(1)} \cdot 1 = h_2 S^{-1}(h_1) = \varepsilon(h) 1.$$

For any object (M, α_M) in ${}_H\mathcal{YD}^H$ and for all $h \in H, m \in M$,

$$\begin{aligned}\rho(h \cdot m) &= \alpha_H^{-1}(h_{21}) \cdot m_{(0)} \otimes [\alpha_H^{-2}(h_{22})\alpha_H^{-1}(m_{(1)})]S^{-1}(h_1) \\ &= \alpha_H^{-1}(h_{12}) \cdot m_{(0)} \otimes [\alpha_H^{-1}(h_2)\alpha_H^{-1}(m_{(1)})]S^{-1}\alpha_H^{-1}(h_{11}) \\ &= h_{(0)} \cdot m_{(0)} \otimes h_{(1)} \cdot m_{(1)}.\end{aligned}$$

That is, (M, α_M) is an object in ${}_H\mathcal{M}(H^{op} \otimes H)^H$. Conversely for any object (M, α_M) in ${}_H\mathcal{M}(H^{op} \otimes H)^H$, by the same computation it is also an object in ${}_H\mathcal{YD}^H$. Thus these two categories are isomorphic and have the same monoidal structure.

The proof is completed. $\square$

Let (H, A, C) and (H', A', C') be two Doi-Hopf data. A morphism $\varphi : (H, A, C) \rightarrow (H', A', C')$ consists of three maps $\theta : H \rightarrow H'$, $\beta : A \rightarrow A'$ and $\gamma : C \rightarrow C'$ which are, respectively a Hom-Hopf algebra map, a Hom-algebra map and a Hom-coalgebra map satisfying

$$\gamma(h \cdot c) = \theta(h) \cdot \gamma(c), \quad (2.5)$$

$$\rho(\beta(a)) = \beta(a_{(0)}) \otimes \theta(a_{(1)}) \quad (2.6)$$

for all $h \in H, c \in C$ and $a \in A$.

Proposition 2.6. *The functor $F : {}_A\mathcal{M}(H)^C \longrightarrow {}_{A'}\mathcal{M}(H)^{C'}$ given by*

$$F(M) = A' \otimes_A M,$$

where the Doi-Hopf module structure on $A' \otimes_A M$ is defined in the following way

$$a' \cdot (b' \otimes m) = \alpha_{A'}(a')b' \otimes \alpha_M(m), \quad (2.7)$$

$$\rho(b' \otimes m) = b'_{(0)} \otimes m_{(0)} \otimes \alpha_{M'}^{-2}(b'_{(1)} \cdot \gamma(m_{(1)})), \quad (2.8)$$

for all $a', b' \in A'$ and $m \in M$.

Proof. Clearly for any object (M, α_M) in ${}_A\mathcal{M}(H)^C$, $A' \otimes_A M$ is a left A' -module and right C' -comodule. Then for all $a', b' \in A'$ and $m \in M$,

$$\begin{aligned}\rho(a' \cdot (b' \otimes m)) &= \rho(\alpha_{A'}(a')b' \otimes \alpha_M(m)) \\ &= \alpha_{A'}(a'_{(0)})b'_{(0)} \otimes \alpha_M(m_{(0)}) \otimes \alpha_{C'}^{-2}(\alpha_{A'}(a'_{(1)})b'_{(1)} \cdot \gamma(\alpha_M(m_{(1)}))) \\ &= a'_{(0)} \cdot (b'_{(0)} \otimes m_{(0)}) \otimes a'_{(1)} \cdot \alpha_{C'}^{-2}(b'_{(1)} \cdot \gamma(m_{(1)})) \\ &= a'_{(0)} \cdot (b' \otimes m)_{(0)} \otimes a'_{(1)} \cdot (b' \otimes m)_{(1)}.\end{aligned}$$

This completes the proof. $\square$

Proposition 2.7. *The functor $G : {}_A\mathcal{M}(H)^{C'} \longrightarrow {}_A\mathcal{M}(H)^C$ given by*

$$G(N') = N' \square_{C'} C,$$

where the Doi-Hopf module structure on $N' \square_{C'} C$ is defined in the following way

$$a \cdot (n' \otimes c) = \beta(a_{(0)}) \cdot n' \otimes a_{(1)} \cdot c, \quad (2.9)$$

$$\rho(n' \otimes c) = \alpha_{N'}(n') \otimes c_1 \otimes \alpha_C^{-1}(c_2), \quad (2.10)$$

for all $a \in A, c \in C$ and $n' \in N'$.

Moreover (F, G) is a pair of adjoint functors.

Proof. First of all, we need to show the action and coaction on $G(N')$ is well defined for any object $N' \in {}_A\mathcal{M}(H)^{C'}$. For all $a \in A, c \in C$ and $n' \in N'$,

$$\begin{aligned} & (\beta(a_{(0)}) \cdot n')_{(0)} \otimes (\beta(a_{(0)}) \cdot n')_{(1)} \otimes \alpha_H(a_{(1)}) \cdot \alpha_C(c) \\ &= \beta(a_{(0)(0)}) \cdot n'_{(0)} \otimes a_{(0)(1)} \cdot n'_{(1)} \otimes \alpha_H(a_{(1)}) \cdot \alpha_C(c) \\ &= \alpha_{A'}(\beta(a_{(0)})) \cdot \alpha_{N'}(n') \otimes a_{(1)1} \cdot \gamma(c_1) \otimes a_{(1)2} \cdot c_2, \end{aligned}$$

thus the action is well defined. And

$$\begin{aligned} & \alpha_{N'}(n')_{(0)} \otimes \alpha_{N'}(n')_{(1)} \otimes \alpha_C(c_1) \otimes \alpha_C^{-1}(c_2) \\ &= \alpha_{N'}(n'_{(0)}) \otimes \alpha_{C'}(n'_{(1)}) \otimes \alpha_C(c_1) \otimes \alpha_C^{-1}(c_2) \\ &= \alpha_{N'}^2(n') \otimes \alpha_{C'}(\gamma(c_1)) \otimes c_{21} \otimes \alpha_C^{-2}(c_{22}) \\ &= \alpha_{N'}^2(n') \otimes \gamma(c_{11}) \otimes c_{12} \otimes \alpha_C^{-1}(c_2), \end{aligned}$$

thus the coaction is well defined. It is straightforward to verify that $G(N')$ is a left A -module and right C -comodule. For the compatibility we have

$$\begin{aligned} \rho(a \cdot (n' \otimes c)) &= \alpha_{A'}(\beta(a_{(0)})) \cdot \alpha_{N'}(n') \otimes a_{(1)1} \cdot c_1 \otimes \alpha_H^{-1}(a_{(1)2}) \cdot \alpha_C^{-1}(c_2) \\ &= \beta(a_{(0)(0)}) \cdot \alpha_{N'}(n') \otimes a_{(0)(1)} \cdot c_1 \otimes a_{(1)} \cdot \alpha_C^{-1}(c_2) \\ &= a_{(0)} \cdot (\alpha_{N'}(n') \otimes c_1) \otimes a_{(1)} \cdot \alpha_C^{-1}(c_2). \end{aligned}$$

Then $G(N')$ is an object in ${}_A\mathcal{M}(H)^C$. Finally we show (F, G) is a pair of adjoint functors. Define the linear maps $\eta_M : M \longrightarrow GF(M)$ by

$$m \mapsto (1 \otimes \alpha_M(m_{(0)})) \otimes m_{(1)},$$

and $\delta_{M'} : FG(M') \longrightarrow M'$ by

$$a' \otimes (m' \otimes c) \mapsto \varepsilon(c) \alpha_{A'}^{-2}(a') \cdot \alpha_{M'}^{-3}(m'),$$

For all $a \in A$ and $m \in M$,

$$\begin{aligned}\eta_M(a \cdot m) &= (1 \otimes \alpha_M(a_{(0)} \cdot m_{(0)})) \otimes a_{(1)} \cdot m_{(1)} \\ &= \beta(\alpha^2(a_{(0)})) \otimes \alpha_M^2(m_{(0)}) \otimes a_{(1)} \cdot m_{(1)} \\ &= a \cdot \eta_M(m),\end{aligned}$$

and

$$\begin{aligned}\rho(\eta_M(m)) &= (1 \otimes \alpha_M^2(m_{(0)})) \otimes m_{(1)1} \otimes \alpha_C^{-1}(m_{(1)2}) \\ &= (1 \otimes \alpha_M(m_{(0)(0)})) \otimes m_{(0)(1)} \otimes m_{(1)} \\ &= \eta_M(m_{(0)}) \otimes m_{(1)}.\end{aligned}$$

Hence η_M is a morphism in ${}_A\mathcal{M}(H)^C$. Similarly we can check that $\delta_{M'}$ is a morphism in ${}_{A'}\mathcal{M}(H)^{C'}$. For all $m' \in M, c \in C$,

$$\begin{aligned}G(\delta_{M'}) \circ \eta_{G(M')}(m' \otimes c) &= G(\delta_{M'})(1 \otimes \alpha_{M'}^2(m') \otimes \alpha_C(c_1) \otimes \alpha_C^{-1}(c_1)) \\ &= m' \otimes c,\end{aligned}$$

and for all $a' \in A', m \in M$,

$$\begin{aligned}\delta_{F(M)} \circ F(\eta_M)(a' \otimes m) &= \delta_{F(M)}(a' \otimes ((1 \otimes \alpha_M(m_{(0)})) \otimes m_{(1)})) \\ &= \alpha_{A'}^{-2}(a') \cdot (1 \otimes \alpha_M^{-2}(m_{(0)})) \varepsilon(m_{(1)}) \\ &= a' \otimes m.\end{aligned}$$

The proof is completed. $\square$

In the rest of this section, we will establish tensor identities involving the pair of adjoint functors (F, G) .

Theorem 2.8. *Let $(id, id, \gamma) : (H, A, C) \longrightarrow (H, A, C')$ be a monoidal morphism of Doi-Hopf data. Then $G(C') = C$.*

Furthermore let $M \in {}_A\mathcal{M}(H)^C$ and $N \in {}_{A'}\mathcal{M}(H)^{C'}$ be arbitrary. If (C, α_C) is a Hom-Hopf algebra, we have the following isomorphism in ${}_A\mathcal{M}(H)^C$

$$M \otimes G(N) \simeq G(F(M) \otimes N). \quad (2.11)$$

If (C, α_C) has a twisted antipode, then

$$G(N) \otimes M \simeq G(N \otimes F(M)). \quad (2.12)$$

Proof. First of all, define linear map $f : C' \square_{C'} C \rightarrow C$, by $f(c' \otimes c) = \varepsilon(c')c$. For all $a \in A$,

$$\begin{aligned} f(a \cdot (c' \otimes c)) &= f(a_{(0)} \cdot c' \otimes a_{(1)} \cdot c) \\ &= f(\varepsilon(a_{(0)(0)})a_{(0)(1)} \cdot c' \otimes a_{(1)} \cdot c) \\ &= \varepsilon(a_{(0)})\varepsilon(c')a_{(1)} \cdot c \\ &= \varepsilon(c')a \cdot c = a \cdot f(c' \otimes c), \end{aligned}$$

that is, f is A -linear. And

$$\begin{aligned} f(c' \otimes c)_{(0)} \otimes f(c' \otimes c)_{(1)} &= \varepsilon(c')c_1 \otimes \alpha_C^{-1}(c_2) \\ &= f((c' \otimes c)_{(0)}) \otimes (c' \otimes c)_{(1)}, \end{aligned}$$

that is, f is C -colinear.

Define $g : C \rightarrow C' \square_{C'} C$ by $g(c) = \gamma \alpha_C^{-1}(c_1) \otimes \alpha_C^{-1}(c_2)$. Easy to see that $g(c) \in C' \square_{C'} C$, and for all $c \in C, c' \in C'$,

$$\begin{aligned} g(f(c' \otimes c)) &= g(\varepsilon(c')c) = \varepsilon(c')\gamma \alpha_C^{-1}(c_1) \otimes \alpha_C^{-1}(c_2) \\ &= \varepsilon(c'_1)\alpha_C^{-1}(c'_2) \otimes c = c' \otimes c, \end{aligned}$$

and

$$f(g(c)) = f(\gamma \alpha_C^{-1}(c_1) \otimes \alpha_C^{-1}(c_2)) = c.$$

Thus f is bijective, and $G(C') = C$.

For any object $M \in {}_A\mathcal{M}(H)^C$ and $N \in {}_A\mathcal{M}(H)^{C'}$, define $\Phi_1 : M \otimes (N \square_{C'} C) \rightarrow (M \otimes N) \square_{C'} C$ by

$$m \otimes (n \otimes c) \mapsto (m_{(0)} \otimes n) \otimes \alpha_C^{-2}(m_{(1)}c),$$

for all $m \in M, n \in N$ and $c \in C$. Easy to see Φ_1 is well defined. We need to show that Φ_1 is a morphism in $M \in {}_A\mathcal{M}(H)^C$. In fact for all $a \in A$,

$$\begin{aligned} \Phi_1(a \cdot (m \otimes (n \otimes c))) &= \Phi_1(a_1 \cdot m \otimes (a_{2(0)} \cdot n \otimes a_{2(1)} \cdot c)) \\ &= (a_{1(0)} \cdot m_{(0)} \otimes a_{2(0)} \cdot n) \otimes \alpha_C^{-2}((a_{1(1)} \cdot m_{(1)})(a_{2(1)} \cdot c)) \\ &\stackrel{(2.1)}{=} (a_{(0)1} \cdot m_{(0)} \otimes a_{(0)2} \cdot n) \otimes a_{(1)} \cdot \alpha_C^{-2}(m_{(1)}c) \\ &= a_{(0)} \cdot (m_{(0)} \otimes n) \otimes a_{(1)} \cdot \alpha_C^{-2}(m_{(1)}c) \\ &= a \cdot \Phi_1(m \otimes (n \otimes c)), \end{aligned}$$

and

$$\begin{aligned} \rho(\Phi_1(m \otimes (n \otimes c))) &= (\alpha_M(m_{(0)}) \otimes \alpha_N(n)) \otimes \alpha_C^{-2}(m_{(1)1}c_1) \otimes \alpha_C^{-3}(m_{(1)2}c_2) \\ &= (m_{(0)(0)} \otimes \alpha_N(n)) \otimes \alpha_C^{-2}(m_{(0)(1)}c_1) \otimes \alpha_C^{-2}(m_{(1)}\alpha_C^{-1}(c_2)) \\ &= \Phi_1(m_{(0)} \otimes (\alpha_N(n) \otimes c_1)) \otimes \alpha_C^{-2}(m_{(1)}\alpha_C^{-1}(c_2)) \\ &= \Phi_1((m \otimes (n \otimes c))_{(0)}) \otimes (m \otimes (n \otimes c))_{(1)}, \end{aligned}$$

as required. Finally we show that Φ_1 is bijective. Define the linear map $\Psi_1 : (M \otimes N) \square_{C'} C \longrightarrow M \otimes (N \square_{C'} C)$ by

$$(m \otimes n) \otimes c \mapsto \alpha_M^{-2}(m_{(0)}) \otimes (n \otimes S_C(\alpha_C^{-2}(m_{(1)}))c),$$

for all $m \in M, n \in N$ and $c \in C$.

Before proceeding we have to verify that Ψ_1 is well defined. For $(m \otimes n) \otimes c \in (M \otimes N) \square_{C'} C$, we have the relation

$$m_{(0)} \otimes n_{(0)} \otimes \alpha_C^{-2}(\gamma(m_{(1)})n_{(1)}) \otimes \alpha_C(c) = \alpha_M(m) \otimes \alpha_N(n) \otimes \gamma(c_1) \otimes c_2. \quad (2.13)$$

Applying $(id \otimes \gamma \otimes id) \circ (id \otimes \Delta_C \circ S_C) \circ \rho_M \circ \alpha_M^{-3}$ to the above identity, we obtain that

$$\begin{aligned} & \alpha_M^{-3}(m_{(0)(0)}) \otimes \gamma S_C \alpha_C^{-3}(m_{(0)(1)2}) \otimes S_C \alpha_C^{-3}(m_{(0)(1)1}) \otimes n_{(0)} \otimes \alpha_C^{-2}(\gamma(m_{(1)})n_{(1)}) \otimes \alpha_C(c) \\ &= \alpha_M^{-2}(m_{(0)}) \otimes \gamma S_C \alpha_C^{-2}(m_{(1)2}) \otimes S_C \alpha_C^{-2}(m_{(1)1}) \otimes \alpha_N(n) \otimes \gamma(c_1) \otimes c_2. \end{aligned}$$

Multiplying the second and the fifth factor, also the third and sixth factor, we obtain that

$$\begin{aligned} & \alpha_M^{-3}(m_{(0)(0)}) \otimes n_{(0)} \otimes \gamma S_C \alpha_C^{-3}(m_{(0)(1)2}) \alpha_C^{-2}(\gamma(m_{(1)})n_{(1)}) \otimes S_C \alpha_C^{-3}(m_{(0)(1)1}) \alpha_C(c) \\ &= \alpha_M^{-2}(m_{(0)}) \otimes \alpha_N(n) \otimes \gamma S_C \alpha_C^{-2}(m_{(1)2}) \gamma(c_1) \otimes S_C \alpha_C^{-2}(m_{(1)1}) c_2. \end{aligned}$$

Thus

$$\begin{aligned} & \alpha_M^{-2}(m_{(0)}) \otimes n_{(0)} \otimes n_{(1)} \otimes S_C \alpha_C^{-1}(m_{(1)}) \alpha_C(c) \\ &= \alpha_M^{-2}(m_{(0)}) \otimes \alpha_N(n) \otimes \gamma S_C \alpha_C^{-2}(m_{(1)2}) \gamma(c_1) \otimes S_C \alpha_C^{-2}(m_{(1)1}) c_2, \end{aligned}$$

which means that $\Psi_1((m \otimes n) \otimes c) \in M \otimes (N \square_{C'} C)$. Hence it is well defined. Now let us check that Ψ_1 and Φ_1 are mutual inverses.

$$\begin{aligned} \Psi_1 \circ \Phi_1(m \otimes (n \otimes c)) &= \Psi_1((m_{(0)} \otimes n) \otimes \alpha_C^{-2}(m_{(1)}c)) \\ &= \alpha_M^{-2}(m_{(0)(0)}) \otimes (n \otimes S_C(\alpha_C^{-2}(m_{(0)(1)}))(\alpha_C^{-2}(m_{(1)}c))) \\ &= \alpha_M^{-1}(m_{(0)}) \otimes (n \otimes S_C(\alpha_C^{-2}(m_{(1)1}))(\alpha_C^{-3}(m_{(1)2})\alpha_C^{-2}(c))) \\ &= m \otimes (n \otimes c), \end{aligned}$$

and

$$\begin{aligned} \Phi_1 \circ \Psi_1((m \otimes n) \otimes c) &= \Psi_1(\alpha_M^{-2}(m_{(0)}) \otimes (n \otimes S_C(\alpha_C^{-2}(m_{(1)}))c)) \\ &= \alpha_M^{-2}(m_{(0)(0)}) \otimes n \otimes \alpha_C^{-2}(\alpha_C^{-2}(m_{(0)(1)}))(S_C(\alpha_C^{-2}(m_{(1)}))c) \\ &= (m \otimes n) \otimes c. \end{aligned}$$

Hence $M \otimes G(N) \simeq G(F(M) \otimes N)$. Similarly one can prove $G(N) \otimes M \simeq G(N \otimes F(M))$.

The proof is completed. $\square$

Corollary 2.9. *Let (H, A, C) be a monoidal Doi-Hopf datum, and $T : {}_A\mathcal{M}(H)^C \longrightarrow {}_H M$ a functor forgetting the right C -coaction. Then for any Doi-Hopf module M we have the isomorphism in ${}_A\mathcal{M}(H)^C$:*

$$M \otimes C \simeq T(M) \otimes C.$$

Proof. We have the following monoidal morphism

$$(id, id, \varepsilon) : (H, A, C) \rightarrow (H, A, k).$$

Obviously ${}_A M(H)^k \simeq {}_A M$, and $F : {}_A \mathcal{M}(H)^C \simeq {}_A M$ is the functor forgetting the C -coaction. Hence

$$\begin{aligned} M \otimes C &\simeq M \otimes G(k) \simeq G(F(M) \otimes k) \\ &= GF(M) = F(M) \otimes C = T(M) \otimes C. \end{aligned}$$

The proof is completed. $\square$

We have the dual version of Theorem 2.8.

Theorem 2.10. *Let $(id, \lambda, id) : (H, A, C) \longrightarrow (H, A', C)$ be a monoidal morphism of Doi-Hopf data. Then $F(A) = A'$.*

For all $M \in {}_A \mathcal{M}(H)^C$ and $N \in {}_{A'} M(H)^C$, suppose that (A, α_A) is a Hom-Hopf algebra, we have the following isomorphism in $M \in {}_{A'} \mathcal{M}(H)^C$

$$F(M) \otimes N \simeq F(M \otimes G(N)). \quad (2.14)$$

If $(A', \alpha_{A'})$ has a twisted antipode,

$$N \otimes F(M) \simeq F(G(N) \otimes M). \quad (2.15)$$

Proof. First of all remind that (A, α_A) is a left module over itself under the action $a \cdot b = \alpha_A^{-1}(a)b$ for all $a, b \in A$.

Define a linear map $f : A' \otimes_A A \rightarrow A$ by $a' \otimes a \mapsto \alpha_{A'}^{-3}(a')\beta(\alpha_A^{-2}(a))$. It is a routine exercise to check that f is well defined and a morphism in ${}_{A'} M(H)^C$. Its inverse is given by $f^{-1} : A' \rightarrow A' \otimes_A A, a' \mapsto \alpha_{A'}^2(a') \otimes 1$. The first assertion is proved.

If $M \in {}_A \mathcal{M}(H)^C$ and $N \in {}_{A'} M(H)^C$, and (A, α_A) is a Hom-Hopf algebra, define $\Phi_2 : A' \otimes_A (M \otimes G(N)) \longrightarrow (A' \otimes_A M) \otimes N$ by

$$a' \otimes (m \otimes n) \mapsto (a'_1 \otimes m) \otimes \alpha_N^{-2}(a'_2 \cdot n),$$

for all $a' \in A', m \in M$ and $n \in N$. We claim that Φ_2 is well defined and a morphism in ${}_{A'} M(H)^C$. Indeed for all $a \in A$,

$$\begin{aligned} \Phi_2(a'\beta(a) \otimes (\alpha_M(m) \otimes \alpha_N(n))) &= (a'_1\beta(a_1) \otimes \alpha_M(m)) \otimes \alpha_N^{-2}(a'_2\beta(a_2) \cdot \alpha_N(n)) \\ &= (\alpha_{A'}(a'_1) \otimes a_1 \cdot m) \otimes \alpha_N^{-2}(a'_2\beta(a_2) \cdot \alpha_N(n)) \\ &= \Phi_2(\alpha_{A'}(a') \otimes (a \cdot (m \otimes n))) \end{aligned}$$

Then for all $b' \in A'$,

$$\begin{aligned}
\Phi_2(a' \cdot (b' \otimes (m \otimes n))) &= \Phi_2(\alpha_{A'}(a')b' \otimes (\alpha_M(m) \otimes \alpha_N(n))) \\
&= (\alpha_{A'}(a'_1)b'_1 \otimes \alpha_M(m)) \otimes \alpha_N^{-2}(\alpha_{A'}(a'_2)b'_2 \cdot \alpha_N(n)) \\
&= a'_1 \cdot (b'_1 \otimes m) \otimes a'_2 \cdot \alpha_N^{-2}(b'_2 \cdot n) \\
&= a' \cdot \Phi_2(b' \otimes (m \otimes n)),
\end{aligned}$$

and

$$\begin{aligned}
&\Phi_2(a' \otimes (m \otimes n))_{(0)} \otimes \Phi_2(a' \otimes (m \otimes n))_{(1)} \\
&= (a'_1 \otimes m)_{(0)} \otimes \alpha_N^{-2}(a'_2 \cdot n)_{(0)} \otimes \alpha_C^{-2}((a'_1 \otimes m)_{(1)} \cdot \alpha_N^{-2}(a'_2 \cdot n)_{(1)}) \\
&= (a'_{1(0)} \otimes m_{(0)}) \otimes \alpha_N^{-2}(a'_{2(0)} \cdot n_{(0)}) \otimes \alpha_C^{-4}(a'_{1(1)} \cdot m_{(1)}) \cdot \alpha_C^{-4}(a'_{2(1)} \cdot n_{(1)}) \\
&= (a'_{(0)1} \otimes m_{(0)}) \otimes \alpha_N^{-2}(a'_{(0)2} \cdot n_{(0)}) \otimes \alpha_C^{-2}(a'_1) \cdot \alpha_C^{-4}(m_{(1)}n_{(1)}) \\
&= \Phi_2((a' \otimes (m \otimes n))_{(0)}) \otimes (a' \otimes (m \otimes n))_{(1)},
\end{aligned}$$

as needed.

For the inverse of Φ_2 , define the linear map $\Psi_2 : (A' \otimes_A M) \otimes N \longrightarrow A' \otimes_A (M \otimes G(N))$

by

$$(a' \otimes m) \otimes n \mapsto \alpha_{A'}^{-2}(a'_1) \otimes (m \otimes S\alpha_{A'}^{-2}(a'_2) \cdot n),$$

for all $a' \in A', m \in M$ and $n \in N$.

Ψ_2 is well defined since for all $a \in A$,

$$\begin{aligned}
&\Psi_2((a' \beta(a) \otimes \alpha_M(m)) \otimes n) \\
&= \alpha_{A'}^{-2}(a'_1 \beta(a_1)) \otimes (\alpha_M(m) \otimes S\alpha_{A'}^{-2}(a'_2 \beta(a_2)) \cdot n) \\
&= \alpha_{A'}^{-1}(a'_1) \otimes \alpha_A^{-2}(a_1) \cdot (m \otimes S\alpha_{A'}^{-3}(a'_2 \beta(a_2)) \cdot \alpha_N^{-1}(n)) \\
&= \alpha_{A'}^{-1}(a'_1) \otimes (\alpha_A^{-2}(a_{11}) \cdot m \otimes \beta(\alpha_A^{-3})(a_{12})[S\alpha_{A'}^{-3}(a'_2 \beta(a_2))]) \cdot n \\
&= \alpha_{A'}^{-1}(a'_1) \otimes (\alpha_A^{-1}(a_1) \cdot m \otimes [\beta(\alpha_A^{-4}(a_{21}))S\alpha_{A'}^{-4}(\beta(a_{22}))])S\alpha_{A'}^{-3}(a'_2) \cdot n \\
&= \alpha_{A'}^{-1}(a'_1) \otimes (a \cdot m \otimes S\alpha_{A'}^{-2}(a'_2) \cdot n) \\
&= \Psi_2((\alpha_{A'}(a') \otimes a \cdot m) \otimes n).
\end{aligned}$$

Then

$$\begin{aligned}
\Phi_2\Psi_2((a' \otimes m) \otimes n) &= \Phi_2(\alpha_{A'}^{-2}(a'_1) \otimes (m \otimes S\alpha_{A'}^{-2}(a'_2) \cdot n)) \\
&= \alpha_{A'}^{-2}(a'_{11}) \otimes m \otimes \alpha_N^{-2}(\alpha_{A'}^{-2}(a'_{12}) \cdot (S\alpha_{A'}^{-2}(a'_2) \cdot n)) \\
&= \alpha_{A'}^{-1}(a'_1) \otimes m \otimes \alpha_N^{-2}(\alpha_{A'}^{-3}(a'_{12})S\alpha_{A'}^{-3}(a'_{22}) \cdot \alpha_N(n)) \\
&= (a' \otimes m) \otimes n,
\end{aligned}$$

and

$$\begin{aligned}
\Psi_2\Phi_2(a' \otimes (m \otimes n)) &= \Psi_2((a'_1 \otimes m) \otimes \alpha_N^{-2}(a'_2 \cdot n)) \\
&= \alpha_{A'}^{-2}(a'_{11}) \otimes m \otimes S\alpha_{A'}^{-2}(a'_{12}) \cdot \alpha_N^{-2}(a'_2 \cdot n) \\
&= a' \otimes (m \otimes n).
\end{aligned}$$

If $(A', \alpha_{A'})$ has a twisted antipode, similarly one can prove that $N \otimes F(M) \simeq F(G(N) \otimes M)$.

The proof is completed. $\square$

3 Braided structure on the category ${}_A\mathcal{M}^C$

Let (H, A, C) be a monoidal Doi-Hopf datum. In this section, we will discuss how to make the category ${}_A\mathcal{M}(H)^C$ to be braided.

Consider a map $Q : C \otimes C \rightarrow A \otimes A$ with twisted convolution inverse R . So we have

$$Q \circ \alpha_C^{\otimes 2} = \alpha_A^{\otimes 2} \circ Q, \quad (3.1)$$

$$Q(c_2, d_2)R(c_1, d_1) = \varepsilon(c)\varepsilon(d)1 \otimes 1, \quad (3.2)$$

for all $c, d \in C$.

When necessary we will introduce the following notation: for all $c, d \in C$,

$$Q(c, d) = q(c, d) = Q^1(c, d) \otimes Q^2(c, d) \in A \otimes A.$$

Consider two Doi-Hopf modules M and N , define $t_{M,N} : M \otimes N \rightarrow N \otimes M$ by

$$t_{M,N}(m \otimes n) = Q(\alpha_C^{-2}(n_{(1)}), \alpha_C^{-2}(m_{(1)}))(\alpha_N^{-2}(n_{(0)}) \otimes \alpha_M^{-2}(m_{(0)})), \quad (3.3)$$

for all $m \in M$ and $n \in N$. Since Q is convolution invertible, $t_{M,N}$ is a bijective.

Lemma 3.1. *Let (H, A, C) be a Doi-Hopf datum. Then $A \otimes C$ becomes a Doi-Hopf module with the structure*

$$\begin{aligned} a \cdot (b \otimes c) &= \alpha_A(a)b \otimes \alpha_C(c), \\ \rho(b \otimes c) &= b_{(0)} \otimes c_1 \otimes \alpha_C^{-2}(b_{(1)} \cdot c_2), \end{aligned}$$

for all $a, b \in A$ and $c \in C$.

Proof. Straightforward. $\square$

Lemma 3.2. *With notations as above, the linear map $t_{M,N}$ is A -linear for all objects $(M, \alpha_M), (N, \alpha_N)$ in ${}_A\mathcal{M}(H)^C$ if and only if*

$$Q(a_{2(1)} \cdot d, a_{1(1)} \cdot c)(\alpha_A(a_{2(0)}) \otimes \alpha_A(a_{1(0)})) = \Delta(\alpha_A^2(a))Q(\alpha_C(d), \alpha_C(c)), \quad (3.4)$$

for all $a, b \in A$ and $c, d \in C$.

Proof. Suppose that $t_{A \otimes C, A \otimes C}$ is A -linear. For convenience we apply the switch map twice on $(A \otimes C) \otimes (A \otimes C)$, and consider $t_{A \otimes C, A \otimes C}$ as the map $t : A \otimes A \otimes C \otimes C \longrightarrow A \otimes A \otimes C \otimes C$ given by

$$\begin{aligned} t(a \otimes b \otimes c \otimes d) \\ = Q(\alpha_C^{-3}(b_{(1)} \cdot d_1), \alpha_C^{-3}(a_{(1)} \cdot c_1))(\alpha_A^{-2}(b_{(0)}) \otimes \alpha_A^{-2}(a_{(0)})) \otimes \alpha_C^{-1}(d_1) \otimes \alpha_C^{-1}(c_1), \end{aligned}$$

for all $a, b \in A$ and $c, d \in C$. t is A -linear, so

$$\begin{aligned} t(a \cdot (1 \otimes 1 \otimes c \otimes d)) \\ = t(\alpha_A^2(a_1) \otimes \alpha_A^2(a_2) \otimes \alpha_C(c) \otimes \alpha_C(d)) \\ = Q(\alpha_H^{-1}(a_{2(1)}) \cdot \alpha_C^{-2}(d_2), \alpha_H^{-1}(a_{1(1)}) \cdot \alpha_C^{-2}(c_2))(a_{2(0)} \otimes a_{1(0)}) \otimes d_1 \otimes c_1, \\ = a \cdot t(1 \otimes 1 \otimes c \otimes d) \\ = \Delta_A(\alpha_A(a))Q(\alpha_C(d_2) \otimes \alpha_C(c_2)) \otimes d_1 \otimes c_1. \end{aligned}$$

Applying $id \otimes id \otimes \varepsilon \otimes \varepsilon$ to both sides, we obtain (3.4).

Conversely suppose that (3.4) holds, and consider Doi-Hopf modules M and N . For all $a \in A, m \in M$ and $n \in N$,

$$\begin{aligned} t_{M,N}(a \cdot (m \otimes n)) &= t_{M,N}(a_1 \cdot m \otimes a_2 \cdot n) \\ &= Q(\alpha_C^{-2}(a_{2(1)} \cdot n_{(1)}), \alpha_C^{-2}(a_{1(1)} \cdot m_{(1)}))(\alpha_N^{-2}(a_{2(0)} \cdot n_{(0)}) \otimes \alpha_M^{-2}(a_{1(0)} \cdot m_{(0)})) \\ &= Q(\alpha_C^{-3}(a_{2(1)} \cdot n_{(1)}), \alpha_C^{-3}(a_{1(1)} \cdot m_{(1)}))(\alpha_A^{-2}(a_{2(0)}) \otimes \alpha_A^{-2}(a_{1(0)})) \cdot (\alpha_N^{-1}(n_{(0)}) \otimes \alpha_M^{-1}(m_{(0)})) \\ &= \Delta_A(\alpha_A^{-1}(a))Q(\alpha_C^{-2}(n_{(1)}), \alpha_C^{-2}(m_{(1)})) \cdot (\alpha_N^{-1}(n_{(0)}) \otimes \alpha_M^{-1}(m_{(0)})) \\ &= a \cdot t_{M,N}(m \otimes n). \end{aligned}$$

The proof is completed. □

Example 3.3. In the above lemma, let $C = k$ and $Q = Q(1, 1)$. If $t_{M,N}$ is A -linear, then $Q\Delta^{op}(a) = \Delta(a)Q$, which means that A is quasi-cocommutative.

Lemma 3.4. With notations as above, the linear map $t_{M,N}$ is C -linear for all objects $(M, \alpha_M), (N, \alpha_N)$ in ${}_A\mathcal{M}(H)^C$ if and only if

$$\begin{aligned} Q(d_1 \otimes c_1) \otimes \alpha^{-3}(c_2 d_2) \\ = Q(\alpha_C^{-1}(d_2), \alpha_C^{-1}(e_2))_{(0)} \otimes \alpha_C^{-4} \circ m_C(Q(\alpha_C^{-1}(d_2), \alpha_C^{-1}(e_2))_{(1)} \cdot (d_1 \otimes c_1)), \end{aligned} \quad (3.5)$$

for all $c, d \in C$. The m_C is the multiplication map of C . For $a \otimes b \in A \otimes A$, we denote

$$(a \otimes b)_{(0)} \otimes (a \otimes b)_{(1)} = a_{(0)} \otimes b_{(0)} \otimes a_{(1)} \otimes b_{(1)} \in A \otimes A \otimes H \otimes H.$$

Proof. Suppose that t (defined in the proof of Lemma 3.2) is C -colinear, then for all $c, d \in C$,

$$\begin{aligned}
& t(1 \otimes 1 \otimes c \otimes d)_{(0)} \otimes t(1 \otimes 1 \otimes c \otimes d)_{(1)} \\
&= Q^1(\alpha_C^{-1}(d_2) \otimes \alpha_C^{-1}(c_2))_{(0)} \otimes Q^2(\alpha_C^{-1}(d_2) \otimes \alpha_C^{-1}(c_2))_{(0)} \otimes \alpha_C^{-1}(d_{11}) \otimes \alpha_C^{-1}(c_{11}) \\
&\otimes \alpha_C^{-4} \circ m_C((Q^1(\alpha_C^{-1}(d_2) \otimes \alpha_C^{-1}(c_2))_{(1)} \otimes Q^2(\alpha_C^{-1}(d_2) \otimes \alpha_C^{-1}(c_2))_{(1)}) \cdot (\alpha_C^{-1}(d_{12}) \otimes \alpha_C^{-1}(c_{12}))) \\
&= t((1 \otimes 1 \otimes c \otimes d)_{(0)}) \otimes (1 \otimes 1 \otimes c \otimes d)_{(1)} \\
&= t(1 \otimes 1 \otimes c_1 \otimes d_1) \otimes \alpha_C^{-3}(c_2 d_2) \\
&= Q(\alpha_C^{-1}(d_{12}), \alpha_C^{-1}(c_{12})) \otimes \alpha_C^{-1}(d_{11}) \otimes \alpha_C^{-1}(c_{11}) \otimes \alpha_C^{-3}(c_2 d_2).
\end{aligned}$$

Applying $id \otimes id \otimes \varepsilon \otimes \varepsilon \otimes id$ to both sides, we obtain (3.5).

Conversely suppose that (3.5) holds, and consider Doi-Hopf modules M and N . For all $m \in M$ and $n \in N$,

$$\begin{aligned}
& t_{M,N}(m \otimes n)_{(0)} \otimes t_{M,N}(m \otimes n)_{(1)} \\
&= Q^1(\alpha_C^{-2}(n_{(1)}), \alpha_C^{-2}(m_{(1)}))_{(0)} \cdot \alpha_N^{-2}(n_{(0)(0)}) \otimes Q^2(\alpha_C^{-2}(n_{(1)}), \alpha_C^{-2}(m_{(1)}))_{(0)} \cdot \alpha_M^{-2}(m_{(0)(0)}) \\
&\otimes \alpha_C^{-2} \circ m_C[(Q^1(\alpha_C^{-2}(n_{(1)}), \alpha_C^{-2}(m_{(1)}))_{(1)} \\
&\quad \otimes Q^2(\alpha_C^{-2}(n_{(1)}), \alpha_C^{-2}(m_{(1)}))_{(1)}) \cdot (\alpha_C^{-2}(n_{(0)(1)}) \otimes \alpha_C^{-2}(m_{(0)(1)}))] \\
&= Q^1(\alpha_C^{-3}(n_{(1)2}), \alpha_C^{-3}(m_{(1)2}))_{(0)} \cdot \alpha_N^{-1}(n_{(0)}) \otimes Q^2(\alpha_C^{-3}(n_{(1)2}), \alpha_C^{-3}(m_{(1)2}))_{(0)} \alpha_M^{-1}(m_{(0)}) \\
&\otimes \alpha_C^{-2} \circ m_C[(Q^1(\alpha_C^{-3}(n_{(1)2}), \alpha_C^{-3}(m_{(1)2}))_{(1)} \\
&\quad \otimes Q^2(\alpha_C^{-3}(n_{(1)2}), \alpha_C^{-3}(m_{(1)2}))_{(1)}) \cdot (\alpha_C^{-2}(n_{(1)1}) \otimes \alpha_C^{-2}(m_{(1)1}))] \\
&\stackrel{(3.5)}{=} Q(\alpha_C^{-2}(n_{(1)1}), \alpha_C^{-2}(m_{(1)1})) \cdot (\alpha_N^{-1}(n_{(0)}) \otimes \alpha_M^{-1}(m_{(0)})) \otimes \alpha_C^{-3}(m_{(1)2} n_{(1)2}) \\
&= t_{M,N}(m_{(0)} \otimes n_{(0)}) \otimes (m \otimes n)_{(1)}.
\end{aligned}$$

The proof is completed. □

Example 3.5. In the above lemma, suppose that $A = k$. If $t_{M,N}$ is C -colinear, then $Q(d_2, c_2)d_1c_1 = Q(d_1, c_1)c_2d_2$, which means that C is quasi-commutative.

Lemma 3.6. With notations as above, the relation (1.1) holds for all objects M, N, P in ${}_A\mathcal{M}(H)^C$ if and only if

$$\begin{aligned}
& (\Delta \otimes \alpha_A)Q(\alpha_C^{-2}(d)\alpha_C^{-1}(e), \alpha_C^{-1}(c)) \\
&= \alpha_A^2 Q^1(\alpha_C^{-2}(d), \alpha_C^{-3}(c_2)) \\
&\quad \otimes q(e, \alpha_H^{-1}(Q^2(\alpha_C^{-2}(d), \alpha_C^{-3}(c_2))_{(1)}) \cdot \alpha_C^{-3}(c_1))(1 \otimes Q^2(\alpha_C^{-2}(d), \alpha_C^{-3}(c_2))_{(0)}), (3.6)
\end{aligned}$$

for all $a, b \in A$ and $c, d \in C$.

Proof. Let M, N, P be Doi-Hopf modules. For all $m \in M, n \in N$ and $p \in P$, we have the left side of (1.1):

$$\begin{aligned}
& a_{N,P,M} t_{M,N \otimes P} a_{M,N,P} ((m \otimes n) \otimes p) \\
&= a_{N,P,M} [Q(\alpha_C^{-2}((n \otimes \alpha_P(p))_{(1)}), \alpha_C^{-3}(m_{(1)})) \cdot \\
&\quad \cdot ((\alpha_N^{-2} \otimes \alpha_P^{-2})((n \otimes \alpha_P(p))_{(0)}) \otimes \alpha_M^{-3}(m_{(0)}))] \\
&= a_{N,P,M} [(\Delta_A \otimes id)Q(\alpha_C^{-4}(n_{(1)}\alpha_P(p_{(1)})), \alpha_C^{-3}(m_{(1)})) \cdot \\
&\quad \cdot (\alpha_N^{-2}(n_{(0)}) \otimes \alpha_P^{-1}(p_{(0)}) \otimes \alpha_M^{-3}(m_{(0)}))] \\
&= (\alpha_A^{-1} \otimes id \otimes id)(\Delta_A \otimes \alpha_A)Q(\alpha_C^{-4}(n_{(1)}\alpha_C(p_{(1)})), \alpha_C^{-3}(m_{(1)})) \cdot \\
&\quad \cdot (\alpha_N^{-3}(n_{(0)}) \otimes \alpha_P^{-1}(p_{(0)}) \otimes \alpha_M^{-2}(m_{(0)})),
\end{aligned}$$

and the right side of (1.1):

$$\begin{aligned}
& (id_N \otimes t_{M,P})a_{N,M,P}(t_{M,N} \otimes id_P)((m \otimes n) \otimes p) \\
&= (id_N \otimes t_{M,P})[\alpha_N^{-3}(Q^1(n_{(1)}, m_{(1)}) \cdot n_{(0)}) \otimes \alpha_A^{-2}(Q^2(n_{(1)}, m_{(1)})) \cdot \alpha_M^{-2}(m_{(0)}) \otimes \alpha_P(p)] \\
&= \alpha_N^{-3}(Q^1(n_{(1)}, m_{(1)}) \cdot n_{(0)}) \otimes q(\alpha_C^{-1}(p_{(1)}), \alpha_H^{-4}(Q^2(n_{(1)}, m_{(1)})_{(1)}) \cdot \alpha_C^{-4}(m_{(0)(1)})) \cdot \\
&\quad \cdot (\alpha_P^{-1}(p_{(0)}) \otimes \alpha_M^{-4}(Q^2(n_{(1)}, m_{(1)})_{(0)} \cdot m_{(0)(0)})).
\end{aligned}$$

Clearly if (3.6) holds, we obtain the relation (1.1).

Conversely suppose that the relation (1.1) holds in the category ${}_A\mathcal{M}(H)^C$. Taking $U = V = W = A \otimes C$ and applying the switch map as in the previous lemma, we regard both sides of (1.1) as maps

$$A^{\otimes 3} \otimes C^{\otimes 3} \longrightarrow A^{\otimes 3} \otimes C^{\otimes 3}.$$

On one hand we have that

$$\begin{aligned}
& a_{V,W,U} t_{U,V \otimes W} a_{U,V,W} (1 \otimes 1 \otimes 1 \otimes c \otimes d \otimes e) \\
&= (\alpha_A^{-1} \otimes id \otimes id)(\Delta \otimes \alpha_A)Q(\alpha_C^{-3}(d_2)\alpha_C^{-2}(e_2), \alpha_C^{-2}(c_2)) \otimes \alpha_C^{-1}(d_1) \otimes e_1 \otimes \alpha_C^{-1}(c_1),
\end{aligned}$$

and on the other hand

$$\begin{aligned}
& (id_V \otimes t_{U,W})a_{V,U,W}(t_{U,V} \otimes id_W)(1 \otimes 1 \otimes 1 \otimes c \otimes d \otimes e) \\
&= \alpha_A(Q^1(\alpha_C^{-3}(d_2), \alpha_C^{-3}(c_2))) \\
&\quad \otimes q(\alpha_C^{-1}(e_2), \alpha_H^{-1}(Q^2(\alpha_C^{-3}(d_2), \alpha_C^{-3}(c_2))_{(1)}) \cdot \alpha_C^{-4}(c_{12}))(1 \otimes Q^2(\alpha_C^{-3}(d_2), \alpha_C^{-3}(c_2))_{(0)}) \\
&\quad \otimes \alpha_C^{-1}(d_1) \otimes e_1 \otimes \alpha_C^{-2}(c_{11}).
\end{aligned}$$

Applying $id^{\otimes 3} \otimes \varepsilon_C^{\otimes 3}$ to both sides, we obtain (3.6).

The proof is completed. □

Lemma 3.7. *With notations as above, the relation (1.2) holds for all objects M, N, P in ${}_A\mathcal{M}(H)^C$ if and only if*

$$\begin{aligned} & (\alpha_A \otimes \Delta_A)Q(\alpha_C^{-1}(e), \alpha_C^{-1}(c)\alpha_C^{-2}(d)) \\ &= q(\alpha_H^{-1}(Q^1(\alpha_C^{-3}(e_2), \alpha_C^{-2}(d))_{(1)}) \cdot \alpha_C^{-3}(e_1), c)(Q^1(\alpha_C^{-3}(e_2), \alpha_C^{-2}(d))_{(0)} \otimes 1) \\ & \otimes \alpha_A^2(Q^2(\alpha_C^{-3}(e_2), \alpha_C^{-2}(d))), \end{aligned} \quad (3.7)$$

for all $c, d, e \in C$.

Proof. The proof is similar to the proof of the above lemma, and left to the reader. $\square$

We summarize our results as follows.

Theorem 3.8. *Let (H, A, C) be a monoidal Doi-Hopf datum, and $Q : C \otimes C \rightarrow A \otimes A$ a twisted convolution invertible map. Then the family of maps $t_{M,N} : M \otimes N \rightarrow N \otimes M$ given by*

$$t_{M,N}(m \otimes n) = Q(\alpha_C^{-2}(n_{(1)}), \alpha_C^{-2}(m_{(1)})) \cdot (\alpha_N^{-2}(n_{(0)}) \otimes \alpha_M^{-2}(m_{(0)})),$$

defines a braiding on the category ${}_A\mathcal{M}(H)^C$ if and only if Q satisfies the relations 3.4–3.7.

Example 3.9. (1) If $C = k$ and denote $Q = Q(1) = Q^1 \otimes Q^2 = q^1 \otimes q^2 \in A \otimes A$. Then $\alpha_A^{\otimes 2}(Q) = Q(1) = Q$, i.e., Q is α_A -invariant.

Relations (3.6) and (3.7) can be rewritten as

$$\begin{aligned} (\Delta_A \otimes \alpha_A)Q &= \alpha_A^2(Q^1) \otimes \alpha_A(q^1) \otimes q^2 \alpha_A(Q^1) = \alpha_A(Q^1) \otimes \alpha_A(q^1) \otimes q^2 Q^2, \\ (\alpha_A \otimes \Delta_A)Q &= q^1 \alpha_A(Q^1) \otimes \alpha_A(q^1) \otimes \alpha_A^2(Q^2) = q^1 Q^1 \otimes \alpha_A(q^2) \otimes \alpha_A(Q^2). \end{aligned}$$

We conclude that the relations of Theorem 3.8 are satisfied if and only if (A, α_A, Q^{-1}) is quasitriangular.

(2) Hom-bialgebra (H, α_H) is called coquasitriangular if there exists a linear map $\sigma : H \otimes H \rightarrow k$ such that for all $a, b, c \in H$

$$\begin{aligned} \sigma(b_2, a_2)b_1a_1 &= \sigma(b_1, a_1)a_2b_2, \\ \sigma(ab, c) &= \sigma(\alpha_H(a), c_1)\sigma(\alpha_H(b), c_2), \\ \sigma(c, ab) &= \sigma(c_1, \alpha_H(a))\sigma(c_2, \alpha_H(b)). \end{aligned}$$

If $A = k$, relations (3.6) and (3.7) take the form

$$\begin{aligned} Q(de, \alpha_C(c)) &= Q(\alpha_C(e), c_1)Q(\alpha_C(d), c_2), \\ Q(\alpha_C(e), cd) &= Q(e_1, \alpha_C(c))Q(\alpha_C(e_2), d). \end{aligned}$$

We conclude that the relations of Theorem 3.8 are satisfied if and only if (C, α_C, Q) is coquasitriangular.

(3) Let (H, α) be a Hom-Hopf algebra with bijective antipode. We have verified that the category of Yetter-Drinfeld modules ${}_H\mathcal{YD}^H \simeq {}_H\mathcal{M}(H^{op} \otimes H)^H$ is a monoidal category. As we all know, ${}_H\mathcal{YD}^H$ is braided and the braiding is given by

$$t_{M,N} : M \otimes N \rightarrow N \otimes M, \quad m \otimes n \mapsto \alpha_N^{-1}(n_{(0)}) \otimes \alpha_M^{-1}(\alpha_H^{-1}(n_{(1)}) \cdot m).$$

The corresponding map Q is

$$Q(h \otimes g) = \varepsilon(g)(1 \otimes h).$$

It is straightforward to check that Q satisfies the conditions in Theorem 3.8.

4 The generalized smash product of Hom-bialgebra and Drinfeld double

In this section, we will introduce a new algebra such that its module category is isomorphic to the category ${}_A\mathcal{M}(H)^C$. Moreover this new algebra is more general than Drinfeld double and is quasitriangular when ${}_A\mathcal{M}(H)^C$ is braided.

Proposition 4.1. *Let (H, α_H) be a Hom-bialgebra, (A, α_A) a right H -comodule algebra and (B, α_B) a left H -module algebra. Then we have a Hom-algebra $(A \# B, \alpha_A \otimes \alpha_B)$ which is equal to $A \otimes B$ as a vector space, with the following multiplication*

$$(a \# b)(a' \# b') = a\alpha_A^{-1}(a'_{(0)}) \# (\alpha_B^{-1}(b) \leftarrow \alpha_H^{-2}(a'_{(1)}))b',$$

for all $a, a' \in A$ and $b, b' \in B$. The unit is $1_A \# 1_B$.

Proof. The proof is a routine computation. □

If (C, α_C) is a left H -module coalgebra, then $(C^*, (\alpha_C^{-1})^*)$ is a right H -module algebra, and the right H -action is given by

$$\langle f \leftarrow h, c \rangle = \langle f, h \cdot \alpha_C^{-2}(c) \rangle,$$

for all $f \in C^*, c \in C$ and $h \in H$.

Lemma 4.2. *Let (H, A, C) be a Hom-type Doi-Hopf datum. Then we have the category isomorphism ${}_A\mathcal{M}(H)^C \simeq {}_{A \# C^*}\mathcal{M}$.*

Proof. For any object (M, α_M) in ${}_A\mathcal{M}(H)^C$, define the left action of $A \# C^*$ on M by

$$(a \# f) \cdot m = \langle f, m_{(1)} \rangle a \cdot \alpha_M^{-1}(m_{(0)}),$$

for all $a \in A, f \in C^*$ and $m \in M$. Indeed for all $b \in A$ and $g \in C^*$,

$$\begin{aligned}
(a \# f)(b \# g) \cdot \alpha_M(m) &= (a \alpha_A^{-1}(b_{(0)}) \# (\alpha^*(f) \leftarrow \alpha_H^{-2}(b_{(1)})) g) \cdot \alpha_M(m) \\
&= \langle (\alpha^*(f) \leftarrow \alpha_H^{-2}(b_{(1)})) g, \alpha_M(m_{(1)}) \rangle a \alpha_A^{-1}(b_{(0)}) \cdot m_{(0)} \\
&= \langle f, \alpha_H^{-1}(b_{(1)}) \cdot \alpha_M^{-2}(m_{(1)1}) \rangle \langle g, \alpha_M^{-1}(m_{(1)2}) \rangle a \alpha_A^{-1}(b_{(0)}) \cdot m_{(0)} \\
&= \langle g, m_{(1)} \rangle \langle f, \alpha_H^{-1}(b_{(1)}) \cdot \alpha_C^{-2}(m_{(0)(1)}) \rangle \\
&\quad \alpha_A(a) \cdot \alpha_M^{-1}(b_{(0)}) \cdot \alpha_M^{-1}(m_{(0)(0)}) \\
&= \langle g, m_{(1)} \rangle (\alpha_A(a) \# (\alpha^{-1})^*(f)) \cdot (b \cdot \alpha^{-1}(m_{(0)})) \\
&= (\alpha_A(a) \# (\alpha^{-1})^*(f)) \cdot ((b \# g) \cdot m),
\end{aligned}$$

and $(1 \# \varepsilon_C) \cdot m = \alpha_M(m)$.

Conversely suppose that $M \in {}_A \# C^* \mathcal{M}$. Easy to see that M is a left A -module by

$$a \cdot m = (a \# \varepsilon_C) \cdot m,$$

for all $a \in A$ and $m \in M$.

Also M is a right C -comodule by

$$m_{(0)} \otimes m_{(1)} = \sum (1 \# e^i) \cdot m \otimes e_i,$$

where $\{e_i\}$ and $\{e^i\}$ is dual basis of C .

In fact for all $f, g \in C^*$,

$$\begin{aligned}
\alpha_M((1 \# e^i) \cdot m) \langle f, e_{i1} \rangle \langle g, e_{i2} \rangle &= (1 \# (\alpha_C^{-1})^*(e^i)) \cdot \alpha_M(m) \langle f * g, \alpha_C^2(e_i) \rangle \\
&= (1 \# (\alpha_C)^*(f * g)) \cdot \alpha_M(m) \\
&= (1 \# f) \cdot ((1 \# (\alpha_C)^*(e^i)) \cdot m) \\
&= (1 \# e^j) \cdot ((1 \# e^i) \cdot m) \langle f, e_j \rangle \langle g, \alpha_C(e_i) \rangle,
\end{aligned}$$

Therefore $\sum_i \alpha_M((1 \# e^i) \cdot m) \otimes e_{i1} \otimes e_{i2} = \sum_{i,j} (1 \# e^j) \cdot ((1 \# e^i) \cdot m) \otimes e_j \otimes \alpha_C(e_i)$. And

$$\sum_i (1 \# e^i) \cdot m \varepsilon_C(e_i) = (1 \# \varepsilon_C) \cdot m = \alpha_M(m).$$

For all $a \in A$, we have

$$\begin{aligned}
a_{(0)} \cdot m_{(0)} \otimes a_{(1)} \cdot m_{(1)} &= \sum_i a_{(0)} \cdot ((1 \# e^i) \cdot m) \otimes a_{(1)} \cdot e_i \\
&= \sum_i ((\alpha_A^{-1}(a_{(0)}) \# \varepsilon_C)(1 \# e^i)) \cdot \alpha_M(m) \otimes a_{(1)} \cdot e_i \\
&= \sum_i (a_{(0)} \# (\alpha^{-1})^*(e^i)) \cdot \alpha_M(m) \otimes a_{(1)} \cdot e_i \\
&= \sum_i (1 \# (\alpha^{-1})^*(e^i)) (a \# \varepsilon_C) \cdot \alpha_M(m) \otimes e_i \\
&= \sum_i (1 \# e^i) \cdot (a \cdot m) \otimes e_i \\
&= (a \cdot m)_{(0)} \otimes (a \cdot m)_{(1)}.
\end{aligned}$$

The proof is completed. $\square$

Proposition 4.3. *Let (H, α_H) be a Hom-bialgebra, (A, α_A) a right H -comodule algebra and (B, α_B) a left H -module algebra. Assume also that A and B are Hom-bialgebra. Then $(A \# B, \alpha_A \otimes \alpha_B)$ is a Hom-bialgebra with the tensor coproduct if and only if*

$$\begin{aligned} \Delta_A(a_{(0)}) \otimes b_1 &\leftarrow \alpha_H^{-1}(a_{(1)1}) \otimes b_2 \leftarrow \alpha_H^{-1}(a_{(1)2}) \\ &= a_{1(0)} \otimes a_{2(0)} \otimes (b_1 \leftarrow \alpha_H^{-1}(a_{1(1)})) \otimes (b_2 \leftarrow \alpha_H^{-1}(a_{2(1)})), \end{aligned} \quad (4.1)$$

$$\varepsilon_A(a_{(0)}) \varepsilon_B(b \leftarrow \alpha_H^{-1}(a_{(1)})) = \varepsilon_A(a) \varepsilon_B(b). \quad (4.2)$$

Moreover if both of A and B are Hom-Hopf algebras, then $A \# B$ is also a Hom-Hopf algebra with the antipode

$$S(a \# b) = (1 \# S_B \alpha_B^{-1}(b))(S_A \alpha_A^{-1}(a) \# 1),$$

for all $a \in A$ and $b \in B$.

Proof. The proof is straightforward and left to the reader. $\square$

Proposition 4.4. *Let (H, A, C) be a monoidal Doi-Hopf datum. Then $(A \# C^*, \alpha_A \otimes (\alpha_C^{-1})^*)$ is a Hom-bialgebra and consequently ${}_A \mathcal{M}(H)^C$ and ${}_{A \# C^*} \mathcal{M}$ are isomorphic as monoidal category.*

Proof. For all $a \in A$, $c, d \in C$ and $f \in C^*$,

$$\begin{aligned} &\Delta_A(a_{(0)}) \langle f_1 \leftarrow \alpha_H^{-1}(a_{(1)1}), c \rangle \langle f_2 \leftarrow \alpha_H^{-1}(a_{(1)2}), d \rangle \\ &= \Delta_A(a_{(0)}) \langle f_1, \alpha_H^{-1}(a_{(1)1}) \cdot \alpha_C^{-2}(c) \rangle \langle f_2, \alpha_H^{-1}(a_{(1)2}) \cdot \alpha_C^{-2}(d) \rangle \\ &\stackrel{(2.1)}{=} a_{(0)1} \otimes a_{(0)2} \langle f, \alpha_H^{-1}(a_{(1)}) \cdot \alpha_C^{-4}(cd) \rangle \\ &= a_{(0)1} \otimes a_{(0)2} \langle f \leftarrow \alpha_H^{-1}(a_{(1)}), \alpha_C^{-2}(cd) \rangle \\ &= a_{(0)1} \otimes a_{(0)2} \langle f_1 \leftarrow \alpha_H^{-1}(a_{(1)1}), c \rangle \langle f_2 \leftarrow \alpha_H^{-1}(a_{(1)2}), d \rangle, \end{aligned}$$

and

$$\varepsilon_A(a_{(0)}) \langle f, \alpha_H^{-1}(a_{(1)1}) \cdot 1_C \rangle = \varepsilon_A(a) \langle f, 1_C \rangle.$$

Thus (4.1) and (4.2) are satisfied.

The proof is completed. $\square$

Example 4.5. *Consider the monoidal Doi-Hopf datum $(H, H, H^{op} \otimes H)$ defined in Proposition 2.5. The algebra $A \# C^*$ is nothing else than the Drinfeld double $(H \bowtie H^*, \alpha_H \otimes$*

$(\alpha^{-1})^*$). Thus we recover the multiplication, comultiplication and antipode by

$$\begin{aligned}(a \bowtie f)(b \bowtie g) &= a\alpha_H^{-2}(b_{12}) \bowtie [S^{-1}\alpha_H^{-3}(b_{11}) \rightharpoonup ((\alpha_H^*)^2)(f) \leftharpoonup \alpha_H^{-3}(b_2)]g, \\ \Delta(h \otimes f) &= h_1 \bowtie f_1 \otimes h_2 \bowtie f_2, \\ S(a \bowtie f) &= (1 \# S_H^* \alpha_H^*(f))(S_H \alpha_H^{-1}(a) \# \varepsilon_H),\end{aligned}$$

for all $a, b \in H$, $f, g \in H^*$.

Now let (H, A, C) be a monoidal Doi-Hopf datum and assume that we have a twisted convolution inverse map $Q : C \otimes C \rightarrow A \otimes A$ satisfying (3.4)–(3.7). Q induces a map

$$\tilde{Q} : k \longrightarrow (A \# C^*) \otimes (A \# C^*),$$

for all $c, d \in C$ and $f, g \in A^*$.

The braiding on ${}_A\mathcal{M}(H)^C$ translates into a braiding on ${}_{A \# C^*}\mathcal{M}$. This means that $A \# C^*$ is quasitriangular and the corresponding $R \in (A \# C^*) \otimes (A \# C^*)$ is just $\tilde{Q}(1)$.

In particular for a finite dimensional Hom-Hopf algebra (H, α) , the Drinfeld double $D(H)$ is quasitriangular.

ACKNOWLEDGEMENTS

This work was supported by the NSF of China (No. 11371088) and the NSF of Jiangsu Province (No. BK2012736).

References

- [1] L. Alvarez-Gaum, C. Gomez and G. Sierra, Quantum group interpretation of some conformal field theories. Phys. Lett. B **220**(1989), 142–152.
- [2] N. Aizawa, H. Sato, q -deformation of the Virasoro algebra with central extension. Phys. Lett. B **256** (1991), 185–190.
- [3] S. Caenepeel, G. Militaru, S. Zhu, Crossed modules and Doi-Hopf modules. Israel J. Math. **100**(1997), 221–247.
- [4] S. Caenepeel, F. Van Oystaeyen, B. Zhou, Making the category of Doi-Hopf modules into a braided monoidal category. Algebras Representation Theory **1**(1998), 75–96.
- [5] Y. Doi, Unifying Hopf modules. J. Algebra **153** (1992), 373–385.
- [6] J. T. Hartwig, D. Larsson and S. D. Silvestrov, Deformations of Lie algebras using σ -derivations. J. Algebra **295** (2006), 314–361.

- [7] C. Kassel, Quantum Group. Grad. Texts in Math., vol. **155**, Springer Berlin, 1995.
- [8] S. Majid, Foundations of quantum group theory. Cambridge university press, 1995.
- [9] A. Makhlouf, F. Panaite, Hom-L-R-smash products, Hom-diagonal crossed products and the Drinfel'd double of a Hom-Hopf algebra. J. Algebra **441**(2015), 314–343.
- [10] A. Makhlouf, S. D. Silvestrov, Hom-algebra structure. J. Gen. Lie Theory Appl. **2**(2008), 52–64.
- [11] A. Makhlouf, S. D. Silvestrov, Hom-algebras and Hom-coalgebras. J. Alg. Appl. **9**(2010), 553–589.
- [12] D. E. Radford, J. Towber, Yetter-Drinfeld categories associated to an arbitrary bialgebra. J. Pure Appl. Algebra **87**(1993), 259–279.
- [13] M. E. Sweedler, Hopf Algebras. New York: Benjamin, 1969.
- [14] D. Yau, Hom-quantum groups: I. Quasi-triangular Hom-bialgebras. J. Phys. A: Math. Theor. **45**(2012), 065203.

Daowei Lu, Department of Mathematics, Southeast University, Nanjing, Jiangsu Province, 210096, P. R. China, e-mail: ludaowei620@126.com.

Shuanhong Wang, Department of Mathematics, Southeast University, Nanjing, Jiangsu Province, 210096, P. R. China.